



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Association of Polygenic Liabilities for Major Depression, Bipolar Disorder, and Schizophrenia With Risk for Depression in the Danish Population

Citation for published version:

Psychiat Genomics Consortium 2019, 'Association of Polygenic Liabilities for Major Depression, Bipolar Disorder, and Schizophrenia With Risk for Depression in the Danish Population', *JAMA Psychiatry*, vol. 76, no. 5, pp. 516-525. <https://doi.org/10.1001/jamapsychiatry.2018.4166>

Digital Object Identifier (DOI):

[10.1001/jamapsychiatry.2018.4166](https://doi.org/10.1001/jamapsychiatry.2018.4166)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

JAMA Psychiatry

Publisher Rights Statement:

This is the authors' peer-reviewed manuscript as accepted for publication.

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Number of words in the abstract: 335
Number of words in the manuscript: 3,207
Number of tables: 3
Number of figures: 2

Associations between polygenic liabilities for major depression, bipolar disorder and schizophrenia, and risk for depression in the Danish population

Revision date: October 17, 2018

Katherine L Musliner, PhD^{1,2}, Preben B. Mortensen, DrMedSc^{1,2,3}, John J. McGrath, PhD^{2,4,5}, Nis P. Suppli, MD PhD^{1,6}, David M. Hougaard, DrMedSc^{1,7}, Jonas Bybjerg-Grauholm, MSE^{1,7}, Marie Bækvad-Hansen, PhD^{1,7}, Ole Andreassen, MD⁸, Carsten B. Pedersen, PhD^{1,2,3}, Marianne G. Pedersen, MSc^{1,2,3}, Ole Mors, MD^{1,9}, Merete Nordentoft, PhD^{1,6}, Anders D Børghlum, MD PhD^{1,10}, Thomas Werge, PhD^{1,11,12}, Esben Agerbo, PhD^{2,3}, for the Bipolar Disorder Working Group of the Psychiatric Genomics Consortium

1. iPSYCH, The Lundbeck Foundation Initiative for Integrated Psychiatric Research
2. NCRR, National Center for Register-based research, Business and Social Sciences, Aarhus University, Denmark
3. CIRRAU – Center for Integrated Register-based Research at Aarhus University, Denmark
4. Queensland Brain Institute, The University of Queensland, St Lucia, QLD, Australia
5. Queensland Centre for Mental Health Research, The Park Centre for Mental Health, Wacol, QLD, Australia
6. Mental Health Centre Copenhagen, Capital Region of Denmark, Copenhagen University Hospital, Denmark
7. Department for Congenital Disorders, Statens Serum Institut, Copenhagen, Denmark
8. The Norwegian Centre for Mental Disorders Research, Division of Mental Health and Addiction, University of Oslo and Oslo University Hospital, Norway
9. Psychosis Research Unit, Aarhus University Hospital, Risskov, Denmark
10. Centre for Integrative Sequencing, Department of Biomedicine and iSEQ, Aarhus University, Denmark
11. Department of Clinical Medicine, University of Copenhagen, Blegdamsvej 3B, DK-2200 Copenhagen N, Denmark
12. Institute of Biological Psychiatry, Copenhagen Mental Health services, Copenhagen, Denmark

Contact information for corresponding author:

Katherine L. Musliner

Address: Fuglesangs Alle 26, Building R, 8210 Aarhus V, Denmark

Email: klm@econ.au.dk

Key Points

Question: Do polygenic liabilities for major depression (MD), bipolar disorder (BD) and schizophrenia (SZ) predict depression in the general population?

Findings: In this case-cohort study including 34,573 individuals representative of the Danish population, each 1 standard deviation increase in polygenic risk for MD was associated with a 30% increase in hazard of depression. Polygenic liabilities for SZ and BD were also associated with increased hazard of depression, but to a lesser extent.

Meaning: Polygenic risk scores trained in samples of prevalent MD cases can predict first depression in the general population.

37

Tweet

- 38 New study from iPSYCH shows risk for depression diagnosis in the general population increases by
- 39 30% per standard deviation increase in polygenic risk

Abstract

Importance: Although the usefulness of polygenic risk scores as a measure of genetic liability for major depression (MD) has been established, their capacity to predict who will develop depression in the general population remains relatively unexplored.

Objective: To evaluate whether polygenic risk scores for MD, bipolar disorder (BD) and schizophrenia (SZ) predict depression in the general population, and explore whether these polygenic liabilities are associated with heterogeneity in age at onset and severity at initial depression diagnosis.

Design, Setting and Participants: Participants were drawn from the Danish iPSYCH2012 case-cohort study: a representative sample drawn from the population of Denmark born between May 1, 1981 and December 31, 2005. Hazard of depression was estimated using Cox regressions modified to accommodate the case-cohort design. Case-only analyses were conducted using linear and multinomial regressions. Data analysis was conducted from February 2017 to June 2018.

Exposure: Polygenic risk scores for MD, BD and SZ trained using the most recent GWAS results from the PGC.

Main Outcome and Measures: First depressive episode (ICD-10 code F32) treated in specialty care. Severity at initial diagnosis was measured using ICD-10 code severity specifications (mild, moderate, severe without psychosis, severe with psychosis) and treatment setting (inpatient, outpatient, emergency).

58 **Results:** The sample was 51% male and included 34,573 individuals (14,799 depression cases) aged
59 10-31 years at censoring. Each standard deviation increase in polygenic liability for MD, BD and SZ
60 was associated with 30% (HR=1.30, 95% CI [1.27, 1.33]), 5% (1.05, [1.02, 1.07]) and 12% (1.12,
61 [1.09, 1.15]) increases in hazard of depression, respectively. Among cases, higher polygenic liability
62 for BP was associated with earlier depression onset ($p=.002$).

63 **Conclusions:** Polygenic liability for MD predicts first depression in the general population, supporting
64 the idea that these scores tap into an underlying liability for developing the disorder. The fact that PRS-
65 BD and PRS-SZ also predicted depression is consistent with prior evidence that these disorders share a
66 degree of common genetic overlap. Variation in polygenic liability may contribute slightly to
67 heterogeneity in clinical presentation, but these effects appear minimal.

68

Introduction

Genes play a moderate role in the etiology of depression, with twin-based heritability estimates ranging from 30-40%¹ and SNP-based heritability estimates ranging from 9-29%.²⁻⁴ Large empirical studies of the genetic architecture of depression indicate that it is polygenic, meaning that the contribution of genetic factors is attributable to small effects of hundreds or thousands of genetic variants spread across the genome.^{3,5}

To date, multiple studies have shown small but statistically significant associations between polygenic risk scores (PRS) – a weighted sum of the number of variants associated with the disorder in a different dataset – and depression.⁶⁻⁹ However, these studies focused on prevalent depression, which is more likely than incident depression to be recurrent or chronic.^{10,11} Since the genome-wide association study (GWAS) data underpinning the PRSs are largely based upon prevalent cases, this could suggest that part of the genetic architecture discovered in GWAS studies is linked to chronicity or recurrence rather than the risk of developing the disorder. Therefore, while the usefulness of polygenic risk scores as a measure of genetic liability for depression has been established, their capacity to predict who will develop depression in the general population remains relatively unexplored.

In addition to heterogeneity in chronicity and recurrence, depression is also characterized by substantial variation in characteristics such as age at onset and symptom severity. Research suggests that this variation may be due at least partially to differences in genetic liability. Family studies demonstrate that individuals with a parental history of major depression (MD) are at increased risk for onset of depression at earlier ages,¹²⁻¹⁵ and recent results from the Psychiatric Genomics Consortium (PGC) showed that polygenic risk for depression had a stronger effect on the odds of early vs. late onset

90 depression.⁴ Research also suggests that individuals with severe MD may have a higher genetic burden
91 than individuals with milder symptoms: A recent GWAS in Han Chinese women found an increased
92 genetic signal among individuals with melancholia,¹⁶ and the PGC reported higher PRSs among severe
93 vs. moderate depression cases.⁴

94 Our primary aim in this study was to evaluate the extent to which polygenic liability is associated with
95 risk for first depressive episode in the general population. As a secondary aim, we examined whether
96 polygenic liability is associated with severity and age at onset at first depression diagnosis. Because
97 prior evidence suggests a possible shared genetic etiology between depression and other psychiatric
98 disorders,^{2,4,17} we also examined the extent to which PRSs for bipolar disorder (BD) and schizophrenia
99 (SZ) affect the risk for developing depression in the general population. To accomplish these aims, we
100 used data from the iPSYCH2012 sample, a unique and powerful dataset that links genetic information
101 with longitudinal phenotype data from Danish national registers.

102 **Methods**

103 *Study design*

104 For a detailed description of the iPSYCH2012 sample, see Pedersen et al.¹⁸ Briefly, The iPSYCH2012
105 sample has a case-cohort design¹⁹, which consists of two parts: a random sample (i.e. ‘subcohort’) of
106 individuals drawn from a specified base population (i.e. ‘full cohort’) and all additional cases from the
107 full cohort that were not selected as part of the subcohort. Like a traditional cohort study, a case-cohort
108 study can obtain accurate estimates of hazard and risk using traditional survival analysis, provided the
109 analyses are modified to address issues related to point and variance estimation caused by

oversampling cases.¹⁹⁻²² For a more detailed description of the case-cohort design see the Supplemental Methods.

In the iPSYCH2012 sample, the subcohort consists of a random sample of 30,000 individuals drawn from the full cohort of all singletons born in Denmark between May 1, 1981 and December 31, 2005 who were alive and living in Denmark on their first birthday and had known mothers (N=1,472,762).¹⁸ The full cohort was identified using information from the Danish Civil Registration system (DCRS).²³ The iPSYCH2012 study includes all individuals from the full cohort who were diagnosed with depression in a psychiatric hospital in Denmark between 1991 and 2012 at the age of 10 or older. Information on psychiatric diagnoses was obtained from the Danish Central Psychiatric Research Register (DCPRR).²⁴ Diagnoses are based on the *ICD-10 Classification of Mental and Behavioural Disorders: Diagnostic Criteria for Research*.²⁵

Sample selection

We began by selecting all members of the subcohort and all cases outside the subcohort whose first depression diagnosis in the DCPRR was F32 – ‘depressive episode’. We focused exclusively on individuals with Danish-born parents in order to reduce heterogeneity in genetic ancestry. The sample was further restricted to individuals who were successfully genotyped as part of the iPSYCH2012 sample, passed quality control (QC) measures, and were alive, living in Denmark and at least 10 years old by the end of follow up (December 31, 2012). Seventy-six individuals were removed from the subcohort because their first depression diagnosis in the register was F33 – ‘recurrent depressive disorder’, and we were thus unable to identify their first depressive episode. The final sample included

130 34,573 individuals: 20,082 members of the subcohort (of whom 308 were depression cases) and 14,491
131 depression cases outside the subcohort (Supplemental Figure S1).

132 This study was approved by the Danish Data Protection Agency. By Danish law, register-based studies
133 do not require informed consent.

134 *Genotyping*

135 For a more detailed description of genotyping and quality control (QC) in the iPSYCH2012 cohort, see
136 Pedersen et al.¹⁸ Briefly, DNA was collected from blood spots drawn as part of routine clinical testing
137 at birth and stored at The Danish Neonatal Screening Biobank.²⁶ Blood spots were located for 80,422
138 of the 86,189 members of the original iPSYCH2012 sample (93.3%). Genotyping was performed at the
139 Broad Institute of Harvard and MIT (Cambridge, MA, USA) using the Infinium PsychChip v1.0 array
140 (Illumina CA, San Diego, USA) according to the manufacturer's protocols.²⁷ QC and imputation (using
141 1000 genomes as reference panel) were conducted using the Ricopili pipeline. As the iPSYCH2012
142 sample is population-based, individuals from the same nuclear family unit were neither purposely
143 sampled nor purposely removed.

144 *Measures*

145 Age at onset (AAO) was operationalized as the individual's age in years at first F32 diagnosis in the
146 DCPRR. Information on severity was obtained from the ICD-10 diagnostic code: Mild (F32.0);
147 Moderate (F32.1); Severe without psychotic features (F32.2); Severe with psychotic features (F32.3).
148 We also examined differences in treatment setting (inpatient, emergency, outpatient) as cases treated in
149 inpatient or emergency settings are likely more severe than cases treated in an outpatient setting.

150 *Polygenic risk scores.* PRSs were calculated using a standard approach,²⁸ in which an LD-pruned
151 discovery dataset is used to identify autosomal SNPs associated with the outcome at varying p-value
152 thresholds ($p(t)$), and then a score is calculated for each individual in a target dataset corresponding to
153 the weighted sum of each subject's allelic burden at that threshold. Polygenic risk scores for MD and
154 SZ were created using the Ricopili process with the most recently published GWAS results from the
155 PGC (not including iPSYCH2012) as discovery datasets.^{4,29} The discovery dataset for BD was
156 comprised of leave-one-out summary statistics provided in advance of the latest GWAS publication
157 from the BP Working Group of the PGC.³⁰ SNPs from the discovery datasets were filtered at $INFO >$
158 0.9 and $MAF > .05$, and the broad MHC-region (chr6: 25-35MB) was removed. Additionally SNPs
159 were only included in the scores if they were reliably genotyped or imputed across all 23 waves of the
160 iPSYCH2012 sample, at $INFO > 0.6$ and $MAF > .01$. Ten PRSs were calculated for each disorder (30
161 total) at the following p-value thresholds: $p < .00000005$; 0.000001 ; 0.0001 ; 0.001 ; 0.01 ; 0.05 ; 0.1 ; 0.2 ;
162 0.5 ; 1.0 (See Supplemental Table S1 for SNP numbers).³¹ PRSs were standardized using the means and
163 standard deviations from the distributions in subcohort members with Danish ancestry ($N=22,182$). As
164 the iPSYCH2012 subcohort is a simple random sample drawn from the base population, the
165 distribution of PRSs in the subcohort approximates the distribution in the Danish-born population.

166 *Statistical analysis*

167 Hazard of depression was estimated using Cox regressions with days since 10th birthday as the
168 timescale. Individuals entered the analysis on their 10th birthday and were censored on the date of their
169 first F32 diagnosis in the DCPRR, death, emigration, or December 31, 2012, whichever came first. The
170 oldest participants were 31 years at diagnosis or censoring. We used robust standard errors and Barlow
171 weighting to account for oversampling of cases^{22,32}. All models were adjusted for sex and the top four

ancestral principal components. Additionally, all models were stratified by birth year (1981-2002) to control for secular trends in diagnostic practices and because fewer bloodspots were retrievable among individuals born in earlier years.

To examine whether polygenic liability has a stronger association with early-onset depression, we conducted separate Cox regressions, in each analysis considering as outcome only depression with onset in a specific age range. Thus, we estimated separately the association of PRSs with hazard of depression with diagnosis at ages 10-15, 16-20, 21-25 and 26-31, respectively. Likewise, we conducted separate Cox regressions considering respectively mild, moderate, severe without psychotic features, severe with psychotic features, inpatient, emergency and outpatient treatment setting at first depression as outcome in each of the analyses. We also examined the associations between PRSs, AAO and severity among cases only. Associations with age at onset in days were estimated using linear regressions. Associations with severity measures were estimated using multinomial logistic regressions. All case-only regressions were adjusted for the first 4 PCs, sex and birth year. Statistical significance was assessed at the bonferroni-corrected alpha level $p < .017$. Analyses were conducted in SAS 9.4 (See Appendix in the Supplemental Materials for SAS code).

Results

Sample characteristics

Table 1 shows sample characteristics. Non-cases were 49% female, which was expected given that the subcohort is a representative sample of the Danish population. MD cases were 68% female, which was also expected given that the typical female to male ratio of depression in the population is 2:1.³³ AAO ranged from 10 to 31 years, with a mean of 19.1 years (SD = 4.1 years). Seventeen percent of the cases

193 were classified as mild, 45% as moderate, 9% as severe, 3% as psychotic and 25% had no ICD-10
194 severity specification. The majority (60%) were treated in an outpatient setting.

195 *Polygenic liability and hazard of depression*

196 For all three polygenic risk scores (PRS-MD, PRS-BD and PRS-SZ), the strength of the association
197 increased as the stringency of the $p(t)$ threshold decreased up to $p(t) < .05$ (Supplemental Table S2 and
198 Supplemental Figures S2-S7). For this reason and to maintain consistency with prior research, we
199 present results from the $p(t) < .05$ thresholds throughout the remainder of the manuscript.

200 Each standard deviation (SD) increase in PRS-MD was associated with a 30% increase in the hazard of
201 depression (95% CI = 1.27-1.33; $p < .0001$). In other words, compared to an individual with average
202 polygenic liability, an individual one SD above the population average had a 30% increased risk of
203 depression diagnosis before age 31. The corresponding values for PRS-BD and PRS-SZ were 5%
204 (1.02-1.07; $p < .0001$) and 12% (1.09-1.15; $p < .0001$), respectively (Table S2). Figure 1 shows the
205 associations between PRS deciles and hazard of depression, with the bottom decile (i.e. the lowest 10%
206 of the PRS distribution) as the reference category. Relative to individuals in the bottom decile,
207 individuals in the top decile of PRS-MD had a hazard ratio of 2.55 (2.28-2.85; $p < .0001$). The
208 corresponding values were 1.22 (1.10-1.36; $p < .0001$) for PRS-BD and 1.49 (1.34-1.66; $p < .0001$) for
209 PRS-SZ (Supplemental Table S3).

210 *AAO*

211 As shown in Figure 2 and Table 2, the hazard of depression per SD increase in PRS-MD was slightly
212 higher for first diagnosis between 16-20 years (1.31, 1.27-1.35) and 21-25 years (1.32, 1.27-1.38),

213 compared to 10-15 years (1.27, 1.22-1.33) or 26-31 years (1.24, 1.16-1.32). The effect of PRS-BD was
214 strongest for diagnosis between ages 10-15 years (1.08, 1.04-1.12) and decreased linearly with age.
215 PRS-SZ was also for diagnosis between 10-15 years (1.17, 1.12-1.22), but lowest for diagnosis
216 between 21-25 years (1.08, 1.04-1.13). Case-only analyses showed small associations between higher
217 polygenic risk scores and earlier age at onset across all scores, but only the association for PRS-BD
218 survived correction for multiple testing (Table 3).

219 *Severity*

220 The effects of PRS-MD, PRS-BP and PRS-SZ were all strongest for psychotic depression (Figure 2,
221 Table 2). Differences by severity were most pronounced for PRS-SZ (1.20, 1.09-1.32) and least
222 pronounced for PRS-MD (1.33, 1.23-1.45). In the case-only analyses, none of the associations were
223 statistically significant; however the association between higher PRS-SZ and increased odds of
224 psychotic depression was suggestive (OR=1.10, 95% CI [1.00- 1.21]; $p = .06$) (Table 3).

225 The effects of PRS-MD, PRS-BD and PRS-SZ were larger for inpatient and emergency treatment, but
226 the differences in effect size were small (Figure 2). PRS-MD was marginally associated with increased
227 odds of emergency treatment in the case-only analyses (OR=1.05, 95% CI [1.01- 1.09]; $p = .02$) (Table
228 3).

229 **Discussion**

230 In this study, we found that polygenic risk scores trained using aggregated results from selected
231 samples of prevalent, often recurrent depression cases contributed meaningfully to risk for first
232 depression in the Danish general population. For each SD increase in polygenic liability, hazard of

233 depression increased by 30%. Relative to individuals in the bottom 10% of the polygenic liability
234 distribution, the hazard of depression was 2.55 times higher among individuals in the top 10%. These
235 results suggest that estimates of genetic liability ascertained using prevalent samples are tapping in to
236 an underlying genetic predisposition for developing depression, not just a predisposition to maintain the
237 disorder once it has been established. Polygenic liability for BD and SZ predicted depression to a lesser
238 extent than PRS-MD, which supports the well-documented finding that these disorders share a degree
239 of common genetic etiology.

240 The effect of polygenic liability on age at onset in this study was much smaller than prior findings from
241 family studies would suggest. This could reflect the fact that PRSs and family background are not
242 entirely overlapping measures of genetic risk. Previous research suggests that some, but not all, of the
243 effect of family history of schizophrenia is mediated by polygenic risk.³⁴ If the same holds true for
244 depression, the larger effects identified in family studies may be attributable to the portion of the family
245 history effect not captured by a PRS, or possibly to increased vigilance for psychiatric disorders among
246 multiplex families. It should also be noted that the oldest members of the iPSYCH2012 cohort were
247 only 31 at the end of follow-up, around the median age of onset for depression.³⁵ As a result, the entire
248 sample could be considered ‘early onset.’ It may be that more pronounced differences exist in the
249 effects of PRS-MD on risk for depression at different points across the lifespan, but that these
250 differences are less apparent when comparing groups of younger individuals.

251 We found little association between polygenic liability and greater severity at initial depression
252 diagnosis, which is inconsistent with recent findings from the PGC⁴. In general, past studies with
253 positive findings in this area have focused on severity measures that relate to illness course, such as

254 number of depressive episodes and chronicity of depressive symptoms³⁶⁻³⁸. It could be that polygenic
255 liability has less of an impact on characteristics of the first depressive episode than it does on
256 characteristics of course.

257 It has been suggested previously that stratifying on the phenotype may be a viable method to increase
258 statistical power for identifying genetic variants associated with depression.³⁹ This method has been
259 used with some success by the CONVERGE consortium, which identified a locus significantly
260 associated with depression in Han Chinese women by selecting for highly severe, recurrent female
261 cases¹⁶. Polygenic risk was also found to be differentially associated with subtypes in autism³¹, bipolar
262 disorder and schizophrenia³⁰. However, for depression, greater success in gene discovery was achieved
263 by increasing sample size at the expense of a carefully defined phenotype.^{4,40} In this vein, the results of
264 the current study indicate that the usefulness of further stratification on the phenotype for gene
265 discovery in depression might be more limited than we may have wished.

266 We found some evidence of genetic heterogeneity among depression cases in terms of polygenic
267 liabilities for BD and SZ: there was a suggestive (although not significant) association between PRS-
268 SZ and depression with psychotic symptoms, which makes intuitive sense. PRS-BD was significantly
269 associated with earlier age at MD onset in the case-only analyses, and the case-cohort analyses in
270 separate age groups suggest that PRS-BD and PRS-SZ may be particularly elevated among individuals
271 diagnosed with MD between the ages of 10-15. These results are consistent with past studies^{41,42} and
272 could suggest that a person's degree of genetic liability for BD or SZ may place them at increased risk
273 for different clinical manifestations of depression. However, psychiatric diagnoses are often unstable
274 over time, and both early age at onset and greater severity/psychotic symptoms are robust risk factors

275 for converting to BD or SZ.⁴³⁻⁴⁵ It is therefore possible that these results reflect the fact that many
276 individuals with BD and SZ receive a depression diagnosis during the early stages of their illness.

277 *Strengths and Limitations*

278 The iPSYCH2012 sample has many strengths, including large sample size, population-based sampling
279 and a uniform case definition. The fact that cases were identified through clinical records rather than
280 selected specifically for research increases the relevance for clinical practice. In addition, the ancestral
281 homogeneity of the Danish population reduces the likelihood of confounding by population
282 stratification.

283 It should be noted, however, that although cases in the iPSYCH2012 sample are representative of
284 individuals who received treatment for depression in psychiatric hospitals, they do not include
285 depression cases who are untreated or only treated by general practitioners^{46,47}. To put this into
286 perspective, currently unpublished results show that the majority (85%) of individuals treated for
287 depression in Denmark are treated first by their primary care doctors, though this proportion was lower
288 in younger age groups.⁴⁸ The cases in the iPSYCH2012 sample therefore represent the severe end of
289 the depression distribution in Denmark, which is both a strength and a limitation. Severe cases are
290 likely enriched for genetic determinants¹⁶, however the results may not generalize to milder forms of
291 depression, and they could be biased towards the null due to misclassification. Additionally, there
292 might be too little variation in severity to assess the effect of polygenic liability on severity in this
293 sample. The analyses may also be subject to selection bias, if specialty-treated depression cases are
294 more likely than untreated or primary-care treated cases to experience recurrent episodes. It is also
295 worth noting that because we focused exclusively on first depression, we did not account for

296 subsequent diagnostic conversions to BD or SZ. Individuals who convert may have different genetic
297 profiles; however there was no way to account for this without conditioning on the future, which can
298 introduce bias.⁴⁹ Further research is needed to investigate the impact of polygenic liability on
299 characteristics of course and outcome, including progression to other psychiatric disorders. Finally, the
300 discovery datasets used to create the polygenic risk scores for MD, BD and SZ had different sample
301 sizes, which impacts their statistical power.⁵⁰

302 *Conclusions*

303 In conclusion, we found that polygenic liability predicts depression in the general Danish population.
304 Polygenic liabilities for BD and SZ also predicted depression, which supports the idea that there is a
305 shared genetic predisposition across these disorders. Heterogeneity in age at onset might be partially
306 attributable to underlying genetic differences among depression cases, but these effects appear
307 minimal.

308

Acknowledgements

Conflict of Interest Disclosures: The authors have no conflicts of interest to disclose.

Funding/support: This study was funded by the Lundbeck Foundation, Denmark, grant # R155-2014-1724. The PGC has received major funding from the US National Institute of Mental Health and the US National Institute of Drug Abuse (U01 MH109528 and U01 MH1095320).

Role of the Funder/Sponsor: the funding sources had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

Additional Contributions: We gratefully acknowledge The Broad Institute for genotyping and the Psychiatric Genomics Consortium (PGC) Major Depression and Schizophrenia Work Groups as well as 23andMe for making summary statistics available. We extend particular thanks to the Bipolar Disorders Working Group of the PGC for making leave-one-out summary statistics available in advance of publication. We also extend sincere thanks to Thomas Damm Als, PhD, Aarhus University in Aarhus, Denmark, for generating the principal component variables, and Jakob Grove, PhD, Aarhus University in Aarhus, Denmark, for generating the polygenic risk scores.

Access to Data and Data Analysis: Dr. Musliner had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

328 Members of the Bipolar Disorder Working Group of the Psychiatric Genomics Consortium:

329	Eli A Stahl 1,2,3	352	Veneri Antilla 14
330	Gerome Breen 4,5	353	Swapnil Awasthi 13
331	Andreas J Forstner 6,7,8,9,10	354	Judith A Badner 34
332	Andrew McQuillin 11	355	Marie Bækvad-Hansen 19,35
333	Stephan Ripke 12,13,14	356	Jack D Barchas 36
334	Vassily Trubetskoy 13	357	Nicholas Bass 11
335	Manuel Mattheisen 15,16,17,18,19	358	Michael Bauer 37
336	Yunpeng Wang 20,21	359	Richard Belliveau 12
337	Jonathan R I Coleman 4,5	360	Sarah E Bergen 38
338	Hélène A Gaspar 4,5	361	Carsten Bøcker Pedersen 19,28,29
339	Christiaan A de Leeuw 22	362	Erlend Bøen 39
340	Stacy Steinberg 23	363	Marco Boks 40
341	Jennifer M Whitehead Pavlides 24	364	James Boocock 41
342	Maciej Trzaskowski 25	365	Monika Budde 42
343	Tune H Pers 3,26	366	William Bunney 43
344	Peter A Holmans 27	367	Margit Burmeister 44
345	Liam Abbott 12	368	Jonas Bybjerg-Grauholm 19,35
346	Esben Agerbo 19,28,29	369	William Byerley 45
347	Huda Akil 30	370	Miquel Casas 46,47,48,49
348	Diego Albani 31	371	Felecia Cerrato 12
349	Ney Alliey-Rodriguez 32	372	Pablo Cervantes 50
350	Thomas D Als 15,16,19	373	Kimberly Chambert 12
351	Adebayo Anjorin 33	374	Alexander W Charney 2

- | | | | | | |
|-----|------------------------|-------------|-----|--------------------------|-------------|
| 375 | Danfeng Chen | 12 | 399 | Christine Fraser | 27 |
| 376 | Claire Churchhouse | 12,14 | 400 | Nelson B Freimer | 71 |
| 377 | Toni-Kim Clarke | 51 | 401 | Louise Frisé | 72,73,74 |
| 378 | William Coryell | 52 | 402 | Katrin Gade | 42,75 |
| 379 | David W Craig | 53 | 403 | Diane Gage | 12 |
| 380 | Cristiana Cruceanu | 50,54 | 404 | Julie Garnham | 76 |
| 381 | David Curtis | 55,56 | 405 | Claudia Giambartolomei | 41 |
| 382 | Piotr M Czerski | 57 | 406 | Marianne Giørtz Pedersen | 19,28,29 |
| 383 | Anders M Dale | 58,59,60,61 | 407 | Jaqueline Goldstein | 12 |
| 384 | Simone de Jong | 4,5 | 408 | Scott D Gordon | 77 |
| 385 | Franziska Degenhardt | 8,9 | 409 | Katherine Gordon-Smith | 78 |
| 386 | Jurgen Del-Favero | 62 | 410 | Elaine K Green | 79 |
| 387 | J Raymond DePaulo | 63 | 411 | Melissa J Green | 80 |
| 388 | Srdjan Djurovic | 64,65 | 412 | Tiffany A Greenwood | 60 |
| 389 | Amanda L Dobbyn | 1,2 | 413 | Jakob Grove | 15,16,19,81 |
| 390 | Ashley Dumont | 12 | 414 | Weihua Guan | 82 |
| 391 | Torbjørn Elvsåshagen | 66,67 | 415 | José Guzman Parra | 83 |
| 392 | Valentina Escott-Price | 27 | 416 | Marian L Hamshere | 27 |
| 393 | Chun Chieh Fan | 61 | 417 | Martin Hautzinger | 84 |
| 394 | Sascha B Fischer | 6,10 | 418 | Urs Heilbronner | 42 |
| 395 | Matthew Flickinger | 68 | 419 | Stefan Herms | 6,8,9,10 |
| 396 | Tatiana M Foroud | 69 | 420 | Maria Hipolito | 85 |
| 397 | Liz Forty | 27 | 421 | Per Hoffmann | 6,8,9,10 |
| 398 | Josef Frank | 70 | 422 | Dominic Holland | 58,86 |

423	Laura Huckins 1,2	447	Pamela B Mahon 63,109
424	Stéphane Jamain 87,88	448	Wolfgang Maier 110
425	Jessica S Johnson 1,2	449	Lina Martinsson 73
426	Anders Juréus 38	450	Steve McCarroll 12,111
427	Radhika Kandaswamy 4	451	Peter McGuffin 4
428	Robert Karlsson 38	452	Melvin G McInnis 112
429	James L Kennedy 89,90,91,92	453	James D McKay 113
430	Sarah Kittel-Schneider 93	454	Helena Medeiros 95
431	Sarah V Knott 78	455	Sarah E Medland 77
432	James A Knowles 94,95	456	Fan Meng 30,112
433	Manolis Kogevinas 96	457	Lili Milani 114
434	Anna C Koller 8,9	458	Grant W Montgomery 25
435	Ralph Kupka 97,98,99	459	Derek W Morris 115,116
436	Catharina Lavebratt 72	460	Thomas W Mühleisen 6,117
437	Jacob Lawrence 100	461	Niamh Mullins 4
438	William B Lawson 85	462	Hoang Nguyen 1,2
439	Markus Leber 101	463	Caroline M Nievergelt 60,118
440	Phil H Lee 12,14,102	464	Annelie Nordin Adolfsson 119
441	Shawn E Levy 103	465	Evaristus A Nwulia 85
442	Jun Z Li 104	466	Claire O'Donovan 76
443	Chunyu Liu 105	467	Loes M Olde Loohuis 71
444	Susanne Lucae 106	468	Anil P S Ori 71
445	Anna Maaser 8,9	469	Lilijana Oruc 120
446	Donald J MacIntyre 107,108	470	Urban Ösby 121

- | | |
|-------------------------------------|-----------------------------------|
| 471 Roy H Perlis 122,123 | 495 Janet L Sobell 137 |
| 472 Amy Perry 78 | 496 Christine S holm Hansen 19,35 |
| 473 Andrea Pfennig 37 | 497 Anne T Spijker 138 |
| 474 James B Potash 63 | 498 David St Clair 139 |
| 475 Shaun M Purcell 2,109 | 499 Michael Steffens 140 |
| 476 Eline J Regeer 124 | 500 John S Strauss 91,141 |
| 477 Andreas Reif 93 | 501 Fabian Streit 70 |
| 478 C line S Reinbold 6,10 | 502 Jana Strohmaier 70 |
| 479 John P Rice 125 | 503 Szabolcs Szelinger 142 |
| 480 Fabio Rivas 83 | 504 Robert C Thompson 112 |
| 481 Margarita Rivera 4,126 | 505 Thorgeir E Thorgeirsson 23 |
| 482 Panos Roussos 1,2,127 | 506 Jens Treutlein 70 |
| 483 Douglas M Ruderfer 128 | 507 Helmut Vedder 143 |
| 484 Euijung Ryu 129 | 508 Weiqing Wang 1,2 |
| 485 Cristina S nchez-Mora 46,47,49 | 509 Stanley J Watson 112 |
| 486 Alan F Schatzberg 130 | 510 Thomas W Weickert 80,133 |
| 487 William A Scheftner 131 | 511 Stephanie H Witt 70 |
| 488 Nicholas J Schork 132 | 512 Simon Xi 144 |
| 489 Cynthia Shannon Weickert 80,133 | 513 Wei Xu 145,146 |
| 490 Tatyana Shehtman 60 | 514 Allan H Young 147 |
| 491 Paul D Shilling 60 | 515 Peter Zandi 148 |
| 492 Engilbert Sigurdsson 134 | 516 Peng Zhang 149 |
| 493 Claire Slaney 76 | 517 Sebastian Zollner 112 |
| 494 Olav B Smeland 58,135,136 | 518 Rolf Adolfsson 119 |

- 519 Ingrid Agartz 17,39,150
- 520 Martin Alda 76,151
- 521 Lena Backlund 73
- 522 Bernhard T Baune 152
- 523 Frank Bellivier 153,154,155,156
- 524 Wade H Berrettini 157
- 525 Joanna M Biernacka 129
- 526 Douglas H R Blackwood 51
- 527 Michael Boehnke 68
- 528 Anders D Børglum 15,16,19
- 529 Aiden Corvin 116
- 530 Nicholas Craddock 27
- 531 Mark J Daly 12,14
- 532 Udo Dannlowski 158
- 533 Tõnu Esko 3,111,114,159
- 534 Bruno Etain 153,155,156,160
- 535 Mark Frye 161
- 536 Janice M Fullerton 133,162
- 537 Elliot S Gershon 32,163
- 538 Michael Gill 116
- 539 Fernando Goes 63
- 540 Maria Grigoriu-Serbanescu 164
- 541 Joanna Hauser 57
- 542 David M Hougaard 19,35
- 543 Christina M Hultman 38
- 544 Ian Jones 27
- 545 Lisa A Jones 78
- 546 René S Kahn 2,40
- 547 George Kirov 27
- 548 Mikael Landén 38,165
- 549 Marion Leboyer 88,153,166
- 550 Cathryn M Lewis 4,5,167
- 551 Qingqin S Li 168
- 552 Jolanta Lissowska 169
- 553 Nicholas G Martin 77,170
- 554 Fermin Mayoral 83
- 555 Susan L McElroy 171
- 556 Andrew M McIntosh 51,172
- 557 Francis J McMahon 173
- 558 Ingrid Melle 174,175
- 559 Andres Metspalu 114,176
- 560 Philip B Mitchell 80
- 561 Gunnar Morken 177,178
- 562 Ole Mors 19,179
- 563 Preben Bo Mortensen 15,19,28,29
- 564 Bertram Müller-Myhsok 54,180,181
- 565 Richard M Myers 103
- 566 Benjamin M Neale 3,12,14

- | | | | |
|-----|--|-----|-------------------------------|
| 567 | Vishwajit Nimgaonkar 182 | 586 | Hreinn Stefansson 23 |
| 568 | Merete Nordentoft 19,183 | 587 | Kari Stefansson 23,194 |
| 569 | Markus M Nöthen 8,9 | 588 | Eystein Stordal 195,196 |
| 570 | Michael C O'Donovan 27 | 589 | Patrick F Sullivan 38,197,198 |
| 571 | Ketil J Oedegaard 184,185 | 590 | Gustavo Turecki 199 |
| 572 | Michael J Owen 27 | 591 | Arne E Vaaler 200 |
| 573 | Sara A Paciga 186 | 592 | Eduard Vieta 201 |
| 574 | Carlos Pato 95,187 | 593 | John B Vincent 141 |
| 575 | Michele T Pato 95 | 594 | Thomas Werge 19,202,203 |
| 576 | Danielle Posthuma 22,188 | 595 | John I Nurnberger 204 |
| 577 | Josep Antoni Ramos-Quiroga 46,47,48,49 | 596 | Naomi R Wray 24,25 |
| 578 | Marta Ribasés 46,47,49 | 597 | Arianna Di Florio 27,198 |
| 579 | Marcella Rietschel 70 | 598 | Howard J Edenberg 205 |
| 580 | Guy A Rouleau 189,190 | 599 | Sven Cichon 6,8,10,117 |
| 581 | Martin Schalling 72 | 600 | Roel A Ophoff 40,41,71 |
| 582 | Peter R Schofield 133,162 | 601 | Laura J Scott 68 |
| 583 | Thomas G Schulze 42,63,70,75,173 | 602 | Ole A Andreassen 135,136 |
| 584 | Alessandro Serretti 191 | 603 | John Kelsoe 60 |
| 585 | Jordan W Smoller 12,192,193 | 604 | Pamela Sklar 1,2 |

- 605
- 606

607

1 Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai, New York, NY, US
- 608

2 Department of Psychiatry, Icahn School of Medicine at Mount Sinai, New York, NY, US
- 609

3 Medical and Population Genetics, Broad Institute, Cambridge, MA, US
- 610

4 MRC Social, Genetic and Developmental Psychiatry Centre, King's College London, London, GB

611 5 NIHR BRC for Mental Health, King's College London, London, GB

612 6 Department of Biomedicine, University of Basel, Basel, CH

613 7 Department of Psychiatry (UPK), University of Basel, Basel, CH

614 8 Institute of Human Genetics, University of Bonn, Bonn, DE

615 9 Life&Brain Center, Department of Genomics, University of Bonn, Bonn, DE

616 10 Institute of Medical Genetics and Pathology, University Hospital Basel, Basel, CH

617 11 Division of Psychiatry, University College London, London, GB

618 12 Stanley Center for Psychiatric Research, Broad Institute, Cambridge, MA, US

619 13 Department of Psychiatry and Psychotherapy, Charité - Universitätsmedizin, Berlin, DE

620 14 Analytic and Translational Genetics Unit, Massachusetts General Hospital, Boston, MA, US

621 15 iSEQ, Center for Integrative Sequencing, Aarhus University, Aarhus, DK

622 16 Department of Biomedicine - Human Genetics, Aarhus University, Aarhus, DK

623 17 Department of Clinical Neuroscience, Centre for Psychiatry Research, Karolinska Institutet,
624 Stockholm, SE

625 18 Department of Psychiatry, Psychosomatics and Psychotherapy, Center of Mental Health, University
626 Hospital Würzburg, Würzburg, DE

627 19 iPSYCH, The Lundbeck Foundation Initiative for Integrative Psychiatric Research, DK

628 20 Institute of Biological Psychiatry, Mental Health Centre Sct. Hans, Copenhagen, DK

629 21 Institute of Clinical Medicine, University of Oslo, Oslo, NO

630 22 Department of Complex Trait Genetics, Center for Neurogenomics and Cognitive Research,
631 Amsterdam Neuroscience, Vrije Universiteit Amsterdam, Amsterdam, NL

632 23 deCODE Genetics / Amgen, Reykjavik, IS

633 24 Queensland Brain Institute, The University of Queensland, Brisbane, QLD, AU

634 25 Institute for Molecular Bioscience, The University of Queensland, Brisbane, QLD, AU

635 26 Division of Endocrinology and Center for Basic and Translational Obesity Research, Boston
636 Children's Hospital, Boston, MA, US

637 27 Medical Research Council Centre for Neuropsychiatric Genetics and Genomics, Division of
638 Psychological Medicine and Clinical Neurosciences, Cardiff University, Cardiff, GB

639 28 NCRR, National Centre for Register-Based Research, Aarhus University, Aarhus, DK

640 29 CIRRAU, Centre for Integrated Register-based Research, Aarhus University, Aarhus, DK

641 30 Molecular & Behavioral Neuroscience Institute, University of Michigan, Ann Arbor, MI, US

642 31 NEUROSCIENCE, Istituto Di Ricerche Farmacologiche Mario Negri, Milano, IT

643 32 Department of Psychiatry and Behavioral Neuroscience, University of Chicago, Chicago, IL, US

644 33 Psychiatry, Berkshire Healthcare NHS Foundation Trust, Bracknell, GB

645 34 Psychiatry, Rush University Medical Center, Chicago, IL, US

646 35 Center for Neonatal Screening, Department for Congenital Disorders, Statens Serum Institut,
647 Copenhagen, DK

648 36 Department of Psychiatry, Weill Cornell Medical College, New York, NY, US

649 37 Department of Psychiatry and Psychotherapy, University Hospital Carl Gustav Carus, Technische
650 Universität Dresden, Dresden, DE

651 38 Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, SE

652 39 Department of Psychiatric Research, Diakonhjemmet Hospital, Oslo, NO

653 40 Psychiatry, UMC Utrecht Hersencentrum Rudolf Magnus, Utrecht, NL

654 41 Human Genetics, University of California Los Angeles, Los Angeles, CA, US

655 42 Institute of Psychiatric Phenomics and Genomics (IPPG), University Hospital, LMU Munich,
656 Munich, DE

657 43 Department of Psychiatry and Human Behavior, University of California, Irvine, Irvine, CA, US

658 44 Molecular & Behavioral Neuroscience Institute and Department of Computational Medicine &
659 Bioinformatics, University of Michigan, Ann Arbor, MI, US

660 45 Psychiatry, University of California San Francisco, San Francisco, CA, US

661 46 Instituto de Salud Carlos III, Biomedical Network Research Centre on Mental Health
662 (CIBERSAM), Madrid, ES

663 47 Department of Psychiatry, Hospital Universitari Vall d'Hebron, Barcelona, ES

664 48 Department of Psychiatry and Forensic Medicine, Universitat Autònoma de Barcelona, Barcelona,
665 ES

666 49 Psychiatric Genetics Unit, Group of Psychiatry Mental Health and Addictions, Vall d'Hebron
667 Research Institut (VHIR), Universitat Autònoma de Barcelona, Barcelona, ES

668 50 Department of Psychiatry, Mood Disorders Program, McGill University Health Center, Montreal,
669 QC, CA

670 51 Division of Psychiatry, University of Edinburgh, Edinburgh, GB

671 52 University of Iowa Hospitals and Clinics, Iowa City, IA, US

672 53 Translational Genomics, USC, Phoenix, AZ, US

673 54 Department of Translational Research in Psychiatry, Max Planck Institute of Psychiatry, Munich,
674 DE

675 55 Centre for Psychiatry, Queen Mary University of London, London, GB

676 56 UCL Genetics Institute, University College London, London, GB

677 57 Department of Psychiatry, Laboratory of Psychiatric Genetics, Poznan University of Medical
678 Sciences, Poznan, PL

679 58 Department of Neurosciences, University of California San Diego, La Jolla, CA, US

680 59 Department of Radiology, University of California San Diego, La Jolla, CA, US

681 60 Department of Psychiatry, University of California San Diego, La Jolla, CA, US

682 61 Department of Cognitive Science, University of California San Diego, La Jolla, CA, US

683 62 Applied Molecular Genomics Unit, VIB Department of Molecular Genetics, University of Antwerp,
684 Antwerp, Belgium

685 63 Department of Psychiatry and Behavioral Sciences, Johns Hopkins University School of Medicine,
686 Baltimore, MD, US

687 64 Department of Medical Genetics, Oslo University Hospital Ullevål, Oslo, NO

688 65 NORMENT, KG Jebsen Centre for Psychosis Research, Department of Clinical Science, University
689 of Bergen, Bergen, NO

690 66 Department of Neurology, Oslo University Hospital, Oslo, NO

691 67 NORMENT, KG Jebsen Centre for Psychosis Research, Oslo University Hospital, Oslo, NO

692 68 Center for Statistical Genetics and Department of Biostatistics, University of Michigan, Ann Arbor,
693 MI, US

694 69 Department of Medical & Molecular Genetics, Indiana University, Indianapolis, IN, US

695 70 Department of Genetic Epidemiology in Psychiatry, Central Institute of Mental Health, Medical
696 Faculty Mannheim, Heidelberg University, Mannheim, DE

697 71 Center for Neurobehavioral Genetics, University of California Los Angeles, Los Angeles, CA, US

698 72 Department of Molecular Medicine and Surgery, Karolinska Institutet and Center for Molecular
699 Medicine, Karolinska University Hospital, Stockholm, SE

700 73 Department of Clinical Neuroscience, Karolinska Institutet and Center for Molecular Medicine,
701 Karolinska University Hospital, Stockholm, SE

702 74 Child and Adolescent Psychiatry Research Center, Stockholm, SE

703 75 Department of Psychiatry and Psychotherapy, University Medical Center Göttingen, Göttingen, DE

704 76 Department of Psychiatry, Dalhousie University, Halifax, NS, CA

705 77 Genetics and Computational Biology, QIMR Berghofer Medical Research Institute, Brisbane, QLD,
706 AU

707 78 Department of Psychological Medicine, University of Worcester, Worcester, GB

708 79 School of Biomedical and Healthcare Sciences, Plymouth University Peninsula Schools of Medicine
709 and Dentistry, Plymouth, GB

710 80 School of Psychiatry, University of New South Wales, Sydney, NSW, AU

711 81 Bioinformatics Research Centre, Aarhus University, Aarhus, DK

712 82 Biostatistics, University of Minnesota System, Minneapolis, MN, US

713 83 Mental Health Department, University Regional Hospital, Biomedicine Institute (IBIMA), Málaga,
714 ES

715 84 Department of Psychology, Eberhard Karls Universität Tübingen, Tübingen, DE

716 85 Department of Psychiatry and Behavioral Sciences, Howard University Hospital, Washington, DC,
717 US

718 86 Center for Multimodal Imaging and Genetics, University of California San Diego, La Jolla, CA, US

719 87 Psychiatrie Translationnelle, Inserm U955, Créteil, FR

720 88 Faculté de Médecine, Université Paris Est, Créteil, FR

721 89 Campbell Family Mental Health Research Institute, Centre for Addiction and Mental Health,
722 Toronto, ON, CA

723 90 Neurogenetics Section, Centre for Addiction and Mental Health, Toronto, ON, CA

724 91 Department of Psychiatry, University of Toronto, Toronto, ON, CA

725 92 Institute of Medical Sciences, University of Toronto, Toronto, ON, CA

726 93 Department of Psychiatry, Psychosomatic Medicine and Psychotherapy, University Hospital
727 Frankfurt, Frankfurt am Main, DE

728 94 Cell Biology, SUNY Downstate Medical Center College of Medicine, Brooklyn, NY, US

729 95 Institute for Genomic Health, SUNY Downstate Medical Center College of Medicine, Brooklyn,
730 NY, US

731 96 Center for Research in Environmental Epidemiology (CREAL), Barcelona, ES

732 97 Psychiatry, Altrecht, Utrecht, NL

733 98 Psychiatry, GGZ inGeest, Amsterdam, NL

734 99 Psychiatry, VU medisch centrum, Amsterdam, NL

735 100 Psychiatry, North East London NHS Foundation Trust, Ilford, GB

736 101 Clinic for Psychiatry and Psychotherapy, University Hospital Cologne, Cologne, DE

737 102 Psychiatric and Neurodevelopmental Genetics Unit, Massachusetts General Hospital, Boston, MA,
738 US

739 103 HudsonAlpha Institute for Biotechnology, Huntsville, AL, US

740 104 Department of Human Genetics, University of Michigan, Ann Arbor, MI, US

741 105 Psychiatry, University of Illinois at Chicago College of Medicine, Chicago, IL, US

742 106 Max Planck Institute of Psychiatry, Munich, DE

743 107 Mental Health, NHS 24, Glasgow, GB

744 108 Division of Psychiatry, Centre for Clinical Brain Sciences, University of Edinburgh, Edinburgh,
745 GB

746 109 Psychiatry, Brigham and Women's Hospital, Boston, MA, US

747 110 Department of Psychiatry and Psychotherapy, University of Bonn, Bonn, DE

748 111 Department of Genetics, Harvard Medical School, Boston, MA, US

749 112 Department of Psychiatry, University of Michigan, Ann Arbor, MI, US

750 113 Genetic Cancer Susceptibility Group, International Agency for Research on Cancer, Lyon, FR

751 114 Estonian Genome Center, University of Tartu, Tartu, EE

752 115 Discipline of Biochemistry, Neuroimaging and Cognitive Genomics (NICOG) Centre, National
753 University of Ireland, Galway, Galway, IE

754 116 Neuropsychiatric Genetics Research Group, Dept of Psychiatry and Trinity Translational Medicine
755 Institute, Trinity College Dublin, Dublin, IE

756 117 Institute of Neuroscience and Medicine (INM-1), Research Centre Jülich, Jülich, DE

757 118 Research/Psychiatry, Veterans Affairs San Diego Healthcare System, San Diego, CA, US

758 119 Department of Clinical Sciences, Psychiatry, Umeå University Medical Faculty, Umeå, SE

759 120 Department of Clinical Psychiatry, Psychiatry Clinic, Clinical Center University of Sarajevo,
760 Sarajevo, BA

761 121 Department of Neurobiology, Care sciences, and Society, Karolinska Institutet and Center for
762 Molecular Medicine, Karolinska University Hospital, Stockholm, SE

763 122 Psychiatry, Harvard Medical School, Boston, MA, US

764 123 Division of Clinical Research, Massachusetts General Hospital, Boston, MA, US

765 124 Outpatient Clinic for Bipolar Disorder, Altrecht, Utrecht, NL

766 125 Department of Psychiatry, Washington University in Saint Louis, Saint Louis, MO, US

767 126 Department of Biochemistry and Molecular Biology II, Institute of Neurosciences, Center for
768 Biomedical Research, University of Granada, Granada, ES

769 127 Department of Neuroscience, Icahn School of Medicine at Mount Sinai, New York, NY, US

770 128 Medicine, Psychiatry, Biomedical Informatics, Vanderbilt University Medical Center, Nashville,
771 TN, US

772 129 Department of Health Sciences Research, Mayo Clinic, Rochester, MN, US

773 130 Psychiatry and Behavioral Sciences, Stanford University School of Medicine, Stanford, CA, US
774 131 Rush University Medical Center, Chicago, IL, US
775 132 Scripps Translational Science Institute, La Jolla, CA, US
776 133 Neuroscience Research Australia, Sydney, NSW, AU
777 134 Faculty of Medicine, Department of Psychiatry, School of Health Sciences, University of Iceland,
778 Reykjavik, IS
779 135 Div Mental Health and Addiction, Oslo University Hospital, Oslo, NO
780 136 NORMENT, University of Oslo, Oslo, NO
781 137 Psychiatry and the Behavioral Sciences, University of Southern California, Los Angeles, CA, US
782 138 Mood Disorders, PsyQ, Rotterdam, NL
783 139 Institute for Medical Sciences, University of Aberdeen, Aberdeen, UK
784 140 Research Division, Federal Institute for Drugs and Medical Devices (BfArM), Bonn, DE
785 141 Centre for Addiction and Mental Health, Toronto, ON, CA
786 142 Neurogenomics, TGen, Los Angeles, AZ, US
787 143 Psychiatry, Psychiatrisches Zentrum Nordbaden, Wiesloch, DE
788 144 Computational Sciences Center of Emphasis, Pfizer Global Research and Development,
789 Cambridge, MA, US
790 145 Department of Biostatistics, Princess Margaret Cancer Centre, Toronto, ON, CA
791 146 Dalla Lana School of Public Health, University of Toronto, Toronto, ON, CA
792 147 Psychological Medicine, Institute of Psychiatry, Psychology & Neuroscience, King's College
793 London, London, GB
794 148 Department of Mental Health, Johns Hopkins University Bloomberg School of Public Health,
795 Baltimore, MD, US
796 149 Institute of Genetic Medicine, Johns Hopkins University School of Medicine, Baltimore, MD, US
797 150 NORMENT, KG Jebsen Centre for Psychosis Research, Division of Mental Health and Addiction,
798 Institute of Clinical Medicine and Diakonhjemmet Hospital, University of Oslo, Oslo, NO

799 151 National Institute of Mental Health, Klecany, CZ

800 152 Discipline of Psychiatry, University of Adelaide, Adelaide, SA, AU

801 153 Department of Psychiatry and Addiction Medicine, Assistance Publique - Hôpitaux de Paris, Paris,
802 FR

803 154 Paris Bipolar and TRD Expert Centres, FondaMental Foundation, Paris, FR

804 155 UMR-S1144 Team 1: Biomarkers of relapse and therapeutic response in addiction and mood
805 disorders, INSERM, Paris, FR

806 156 Psychiatry, Université Paris Diderot, Paris, FR

807 157 Psychiatry, University of Pennsylvania, Philadelphia, PA, US

808 158 Department of Psychiatry, University of Münster, Münster, DE

809 159 Division of Endocrinology, Children's Hospital Boston, Boston, MA, US

810 160 Centre for Affective Disorders, Institute of Psychiatry, Psychology and Neuroscience, London, GB

811 161 Department of Psychiatry & Psychology, Mayo Clinic, Rochester, MN, US

812 162 School of Medical Sciences, University of New South Wales, Sydney, NSW, AU

813 163 Department of Human Genetics, University of Chicago, Chicago, IL, US

814 164 Biometric Psychiatric Genetics Research Unit, Alexandru Obregia Clinical Psychiatric Hospital,
815 Bucharest, RO

816 165 Institute of Neuroscience and Physiology, University of Gothenburg, Gothenburg, SE

817 166 INSERM, Paris, FR

818 167 Department of Medical & Molecular Genetics, King's College London, London, GB

819 168 Neuroscience Therapeutic Area, Janssen Research and Development, LLC, Titusville, NJ, US

820 169 Cancer Epidemiology and Prevention, M. Sklodowska-Curie Cancer Center and Institute of
821 Oncology, Warsaw, PL

822 170 School of Psychology, The University of Queensland, Brisbane, QLD, AU

823 171 Research Institute, Lindner Center of HOPE, Mason, OH, US

824 172 Centre for Cognitive Ageing and Cognitive Epidemiology, University of Edinburgh, Edinburgh,
825 GB

826 173 Human Genetics Branch, Intramural Research Program, National Institute of Mental Health,
827 Bethesda, MD, US

828 174 Division of Mental Health and Addiction, Oslo University Hospital, Oslo, NO

829 175 Division of Mental Health and Addiction, University of Oslo, Institute of Clinical Medicine, Oslo,
830 NO

831 176 Institute of Molecular and Cell Biology, University of Tartu, Tartu, EE

832 177 Mental Health, Faculty of Medicine and Health Sciences, Norwegian University of Science and
833 Technology - NTNU, Trondheim, NO

834 178 Psychiatry, St Olavs University Hospital, Trondheim, NO

835 179 Psychosis Research Unit, Aarhus University Hospital, Risskov, DK

836 180 Munich Cluster for Systems Neurology (SyNergy), Munich, DE

837 181 University of Liverpool, Liverpool, GB

838 182 Psychiatry and Human Genetics, University of Pittsburgh, Pittsburgh, PA, US

839 183 Mental Health Services in the Capital Region of Denmark, Mental Health Center Copenhagen,
840 University of Copenhagen, Copenhagen, DK

841 184 Division of Psychiatry, Haukeland Universitetssjukehus, Bergen, NO

842 185 Faculty of Medicine and Dentistry, University of Bergen, Bergen, NO

843 186 Human Genetics and Computational Biomedicine, Pfizer Global Research and Development,
844 Groton, CT, US

845 187 College of Medicine Institute for Genomic Health, SUNY Downstate Medical Center College of
846 Medicine, Brooklyn, NY, US

847 188 Department of Clinical Genetics, Amsterdam Neuroscience, Vrije Universiteit Medical Center,
848 Amsterdam, NL

849 189 Department of Neurology and Neurosurgery, McGill University, Faculty of Medicine, Montreal,
850 QC, CA

851 190 Montreal Neurological Institute and Hospital, Montreal, QC, CA

852 191 Department of Biomedical and NeuroMotor Sciences, University of Bologna, Bologna, IT

853 192 Department of Psychiatry, Massachusetts General Hospital, Boston, MA, US

854 193 Psychiatric and Neurodevelopmental Genetics Unit (PNGU), Massachusetts General Hospital,
855 Boston, MA, US

856 194 Faculty of Medicine, University of Iceland, Reykjavik, IS

857 195 Department of Psychiatry, Hospital Namsos, Namsos, NO

858 196 Department of Neuroscience, Norges Teknisk Naturvitenskapelige Universitet Fakultet for
859 naturvitenskap og teknologi, Trondheim, NO

860 197 Department of Genetics, University of North Carolina at Chapel Hill, Chapel Hill, NC, US

861 198 Department of Psychiatry, University of North Carolina at Chapel Hill, Chapel Hill, NC, US

862 199 Department of Psychiatry, McGill University, Montreal, QC, CA

863 200 Dept of Psychiatry, Sankt Olavs Hospital Universitetssykehuset i Trondheim, Trondheim, NO

864 201 Clinical Institute of Neuroscience, Hospital Clinic, University of Barcelona, IDIBAPS,
865 CIBERSAM, Barcelona, ES

866 202 Institute of Biological Psychiatry, MHC Sct. Hans, Mental Health Services Copenhagen, Roskilde,
867 DK

868 203 Department of Clinical Medicine, University of Copenhagen, Copenhagen, DK

869 204 Psychiatry, Indiana University School of Medicine, Indianapolis, IN, US

870 205 Biochemistry and Molecular Biology, Indiana University School of Medicine, Indianapolis, IN, US

871

872

References

1. Sullivan PF, Neale MC, Kendler KS. Genetic epidemiology of major depression: review and meta-analysis. *Am J Psychiatry*. 2000;157(10):1552-1562.
2. Cross-Disorder Group of the Psychiatric Genomics C, Lee SH, Ripke S, et al. Genetic relationship between five psychiatric disorders estimated from genome-wide SNPs. *Nat Genet*. 2013;45(9):984-994.
3. Peterson RE, Cai N, Bigdeli TB, et al. The Genetic Architecture of Major Depressive Disorder in Han Chinese Women. *JAMA Psychiatry*. 2017;74(2):162-168.
4. Wray NR, Ripke S, Mattheisen M, et al. Genome-wide association analyses identify 44 risk variants and refine the genetic architecture of major depression. *Nat Genet*. 2018;50(5):668-681.
5. Sullivan PF, Daly MJ, O'Donovan M. Genetic architectures of psychiatric disorders: the emerging picture and its implications. *Nat Rev Genet*. 2012;13(8):537-551.
6. Peyrot WJ, Milaneschi Y, Abdellaoui A, et al. Effect of polygenic risk scores on depression in childhood trauma. *Br J Psychiatry*. 2014;205(2):113-119.
7. Chang SC, Glymour MM, Walter S, et al. Genome-wide polygenic scoring for a 14-year long-term average depression phenotype. *Brain Behav*. 2014;4(2):298-311.
8. Musliner KL, Seifuddin F, Judy JA, Pirooznia M, Goes FS, Zandi PP. Polygenic risk, stressful life events and depressive symptoms in older adults: a polygenic score analysis. *Psychol Med*. 2015;45(8):1709-1720.
9. Major Depressive Disorder Working Group of the Psychiatric GC, Ripke S, Wray NR, et al. A mega-analysis of genome-wide association studies for major depressive disorder. *Mol Psychiatry*. 2013;18(4):497-511.
10. Cohen P, Cohen J. The clinician's illusion. *Arch Gen Psychiatry*. 1984;41(12):1178-1182.
11. Waltoft BL, Pedersen CB, Nyegaard M, Hobolth A. The importance of distinguishing between the odds ratio and the incidence rate ratio in GWAS. *BMC Med Genet*. 2015;16:71.
12. Weissman MM, Wickramaratne P, Gameroff MJ, et al. Offspring of Depressed Parents: 30 Years Later. *Am J Psychiatry*. 2016;173(10):1024-1032.

- 902 13. Weissman MM, Wickramaratne P, Merikangas KR, et al. Onset of major depression in early
903 adulthood. Increased familial loading and specificity. *Arch Gen Psychiatry*. 1984;41(12):1136-
904 1143.
- 905 14. Lieb R, Isensee B, Hofler M, Wittchen HU. Parental depression and depression in offspring:
906 evidence for familial characteristics and subtypes? *J Psychiatr Res*. 2002;36(4):237-246.
- 907 15. Musliner KL, Trøjberg BB, Waltoft BL, et al. Parental history of psychiatric diagnoses and
908 unipolar depression: a Danish National Register-based cohort study. *Psychol Med*.
909 2015;45(13):2781-2791.
- 910 16. CONVERGE Consortium. Sparse whole-genome sequencing identifies two loci for major
911 depressive disorder. *Nature*. 2015;523(7562):588-591.
- 912 17. Pettersson E, Larsson H, Lichtenstein P. Common psychiatric disorders share the same genetic
913 origin: a multivariate sibling study of the Swedish population. *Mol Psychiatry*. 2016;21(5):717-
914 721.
- 915 18. Pedersen CB, Bybjerg-Grauholm J, Pedersen MG, et al. The iPSYCH2012 case-cohort sample:
916 new directions for unravelling genetic and environmental architectures of severe mental
917 disorders. *Mol Psychiatry*. 2018;23(1):6-14.
- 918 19. Prentice RL. A case-cohort design for epidemiologic cohort studies and disease prevention
919 trials. *Biometrika*. 1986;73(1):1-11.
- 920 20. Petersen L, Sorensen TI, Andersen PK. Comparison of case-cohort estimators based on data on
921 premature death of adult adoptees. *Stat Med*. 2003;22(24):3795-3803.
- 922 21. Self SG, Prentice RL. Asymptotic distribution theory and efficiency results for case-cohort
923 studies. *The Annals of Statistics*. 1988;16(1):64-81.
- 924 22. Barlow WE. Robust variance estimation for the case-cohort design. *Biometrics*.
925 1994;50(4):1064-1072.
- 926 23. Pedersen CB. The Danish Civil Registration System. *Scand J Public Health*. 2011;39(7
927 Suppl):22-25.
- 928 24. Mors O, Perto GP, Mortensen PB. The Danish Psychiatric Central Research Register. *Scand J*
929 *Public Health*. 2011;39(7 Suppl):54-57.

- 930 25. WHO. *The ICD-10 classification of mental and behavioral disorders: Diagnostic criteria for*
931 *research*. Geneva, Switzerland 1993.
- 932 26. Norgaard-Pedersen B, Hougaard DM. Storage policies and use of the Danish Newborn
933 Screening Biobank. *J Inherit Metab Dis*. 2007;30(4):530-536.
- 934 27. Gunderson KL, Steemers FJ, Ren H, et al. Whole-genome genotyping. *Methods Enzymol*.
935 2006;410:359-376.
- 936 28. International Schizophrenia C, Purcell SM, Wray NR, et al. Common polygenic variation
937 contributes to risk of schizophrenia and bipolar disorder. *Nature*. 2009;460(7256):748-752.
- 938 29. Schizophrenia Working Group of the Psychiatric Genomics C. Biological insights from 108
939 schizophrenia-associated genetic loci. *Nature*. 2014;511(7510):421-427.
- 940 30. Bipolar D, Schizophrenia Working Group of the Psychiatric Genomics Consortium. Electronic
941 address drive, Bipolar D, Schizophrenia Working Group of the Psychiatric Genomics C.
942 Genomic Dissection of Bipolar Disorder and Schizophrenia, Including 28 Subphenotypes. *Cell*.
943 2018;173(7):1705-1715 e1716.
- 944 31. Grove J, Ripke S, Als TD, et al. Common risk variants identified in autism spectrum disorder.
945 *Biorxiv*. 2018.
- 946 32. Barlow WE, Ichikawa L, Rosner D, Izumi S. Analysis of case-cohort designs. *J Clin Epidemiol*.
947 1999;52(12):1165-1172.
- 948 33. Pedersen CB, Mors O, Bertelsen A, et al. A comprehensive nationwide study of the incidence
949 rate and lifetime risk for treated mental disorders. *JAMA Psychiatry*. 2014;71(5):573-581.
- 950 34. Agerbo E, Sullivan PF, Vilhjalmsen BJ, et al. Polygenic Risk Score, Parental Socioeconomic
951 Status, Family History of Psychiatric Disorders, and the Risk for Schizophrenia: A Danish
952 Population-Based Study and Meta-analysis. *JAMA Psychiatry*. 2015;72(7):635-641.
- 953 35. Kessler RC, Berglund P, Demler O, Jin R, Merikangas KR, Walters EE. Lifetime prevalence
954 and age-of-onset distributions of DSM-IV disorders in the National Comorbidity Survey
955 Replication. *Arch Gen Psychiatry*. 2005;62(6):593-602.
- 956 36. Levine ME, Crimmins EM, Prescott CA, Phillips D, Arpawong TE, Lee J. A polygenic risk
957 score associated with measures of depressive symptoms among older adults. *Biodemography*
958 *Soc Biol*. 2014;60(2):199-211.

- 959 37. Mistry S, Harrison JR, Smith DJ, Escott-Price V, Zammit S. The use of polygenic risk scores to
960 identify phenotypes associated with genetic risk of bipolar disorder and depression: A
961 systematic review. *J Affect Disord.* 2018;234:148-155.
- 962 38. Ferentinos P, Rivera M, Ising M, et al. Investigating the genetic variation underlying episodicity
963 in major depressive disorder: suggestive evidence for a bipolar contribution. *J Affect Disord.*
964 2014;155:81-89.
- 965 39. Levinson DF, Mostafavi S, Milaneschi Y, et al. Genetic studies of major depressive disorder:
966 why are there no genome-wide association study findings and what can we do about it? *Biol*
967 *Psychiatry.* 2014;76(7):510-512.
- 968 40. Hyde CL, Nagle MW, Tian C, et al. Identification of 15 genetic loci associated with risk of
969 major depression in individuals of European descent. *Nat Genet.* 2016;48(9):1031-1036.
- 970 41. Verduijn J, Milaneschi Y, Peyrot WJ, et al. Using Clinical Characteristics to Identify Which
971 Patients With Major Depressive Disorder Have a Higher Genetic Load for Three Psychiatric
972 Disorders. *Biol Psychiatry.* 2017;81(4):316-324.
- 973 42. Power RA, Tansey KE, Buttenschon HN, et al. Genome-wide Association for Major Depression
974 Through Age at Onset Stratification: Major Depressive Disorder Working Group of the
975 Psychiatric Genomics Consortium. *Biol Psychiatry.* 2017;81(4):325-335.
- 976 43. Musliner KL, Munk-Olsen T, Mors O, Ostergaard SD. Progression from unipolar depression to
977 schizophrenia. *Acta Psychiatr Scand.* 2017;135(1):42-50.
- 978 44. Musliner KL, Ostergaard SD. Patterns and predictors of conversion to bipolar disorder in 91
979 587 individuals diagnosed with unipolar depression. *Acta Psychiatr Scand.* 2018;137(5):422-
980 432.
- 981 45. Ratheesh A, Davey C, Hetrick S, et al. A systematic review and meta-analysis of prospective
982 transition from major depression to bipolar disorder. *Acta Psychiatr Scand.* 2017;135(4):273-
983 284.
- 984 46. Kuramoto-Crawford SJ, Han B, Jacobus-Kantor L, Mojtabai R. Receipt of Depression
985 Treatment From General Medical Providers and Specialty Mental Health Providers. *Psychiatr*
986 *Serv.* 2016;67(7):758-765.
- 987 47. Thornicroft G, Chatterji S, Evans-Lacko S, et al. Undertreatment of people with major
988 depressive disorder in 21 countries. *Br J Psychiatry.* 2017;210(2):119-124.

989 48. Musliner KL, Liu X, Gasse C, Christensen KS, Bottger MTW, Munk-Olsen T. Incidence of
990 treated depression in primary and specialty care: A comprehensive overview based on Danish
991 population registers. in prep.

992 49. Andersen PK, Keiding N. Interpretability and importance of functionals in competing risks and
993 multistate models. *Stat Med.* 2012;31(11-12):1074-1088.

994 50. Dudbridge F. Power and predictive accuracy of polygenic risk scores. *PLoS Genet.*
995 2013;9(3):e1003348.

996

997

998 Table 1

999 Sample characteristics

		Case status			
		Non-case		Case	
		N	%	N	%
Sex					
	Male	10,115	51.15	4,743	32.05
	Female	9,659	48.85	10,056	67.95
Age at depression diagnosis					
	Non-case	19,774	100.00	-	-
	10-15 years	-	-	2,965	20.04
	16-20 years	-	-	6,653	44.96
	21-25 years	-	-	3,981	26.90
	26-31 years	-	-	1,200	8.11
ICD-10 severity specifier					
	Non-case	19,774	100.00	-	-
	Mild	-	-	2,573	17.39
	Moderate	-	-	6,635	44.83
	Severe without psychotic symptoms	-	-	1,386	9.37
	Severe with psychotic symptoms	-	-	481	3.25
	Severity unspecified	-	-	3,724	25.16
Treatment setting					
	Non-case	19,774	100.00	-	-
	Outpatient	-	-	8,880	60.00
	Inpatient	-	-	2,269	15.33
	Emergency	-	-	3,650	24.66
Total sample		19,774	100.00	14,799	100.00

1000

1001

1002 Table 2

1003 *Associations between PRS-MD, PRS-BD and PRS-SZ and hazard of depression within different age at*
 1004 *onset and severity subgroups*

		<i>MD</i>		<i>BD</i>		<i>SZ</i>	
		<i>HR (95% CI)</i>	<i>p value</i>	<i>HR (95% CI)</i>	<i>p value</i>	<i>HR (95% CI)</i>	<i>p value</i>
Age at depression diagnosis	10-15	1.27 (1.22, 1.33)	<.0001	1.08 (1.04, 1.12)	0.0002	1.17 (1.12, 1.22)	<.0001
	16-20	1.31 (1.27, 1.35)	<.0001	1.04 (1.01, 1.07)	0.008	1.12 (1.09, 1.16)	<.0001
	21-25	1.32 (1.27, 1.38)	<.0001	1.04 (1.00, 1.08)	0.05	1.08 (1.04, 1.13)	0.0001
	26-31	1.24 (1.16, 1.32)	<.0001	1.02 (0.95, 1.10)	0.55	1.12 (1.04, 1.19)	0.0018
ICD-10 severity specifier	Mild	1.31 (1.26, 1.37)	<.0001	1.06 (1.02, 1.11)	0.005	1.09 (1.04, 1.14)	0.0002
	Moderate	1.31 (1.27, 1.35)	<.0001	1.03 (1.00, 1.06)	0.04	1.13 (1.10, 1.17)	<.0001
	Severe	1.24 (1.18, 1.32)	<.0001	1.08 (1.02, 1.14)	0.009	1.11 (1.05, 1.18)	0.0004
	Psychotic	1.33 (1.23, 1.45)	<.0001	1.13 (1.03, 1.24)	0.01	1.20 (1.09, 1.32)	<.0001
	Unspecified	1.30 (1.25, 1.35)	<.0001	1.04 (1.01, 1.08)	0.02	1.11 (1.07, 1.15)	<.0001
Treatment setting	Emergency	1.34 (1.29, 1.39)	<.0001	1.07 (1.03, 1.11)	0.001	1.14 (1.10, 1.18)	<.0001
	Inpatient	1.33 (1.27, 1.39)	<.0001	1.06 (1.01, 1.11)	0.01	1.13 (1.08, 1.18)	<.0001
	Outpatient	1.28 (1.24, 1.32)	<.0001	1.04 (1.01, 1.07)	0.008	1.11 (1.08, 1.14)	<.0001

1005

1006 MD = major depression, BD = bipolar disorder, SZ = schizophrenia, HR = hazard ratio, 95% CI = 95%
 1007 confidence interval.

1008

1009 Table 3

1010 *Case-only analyses of the effects of PRS-MD, PRS-BD and PRS-SZ on age at onset and severity at first depression diagnosis*

		MD				BD				SZ			
		β	SE	OR (95% CI)	p	β	SE	OR (95% CI)	p	β	SE	OR (95% CI)	p
Age at depression diagnosis		-0.05	0.02	.	0.04	-0.07	0.02	.	0.002	-0.05	0.02	.	0.04
ICD-10 severity specifier		----- reference category -----											
	Mild												
	Moderate	-0.00	0.02	1.00 (0.95,1.05)	0.99	-0.03	0.02	0.97 (0.93,1.02)	0.24	0.04	0.02	1.04 (0.99,1.09)	0.08
	Severe	-0.05	0.03	0.95 (0.89,1.01)	0.13	0.02	0.03	1.02 (0.95,1.08)	0.62	0.02	0.03	1.02 (0.95,1.09)	0.58
	Psychotic	0.02	0.05	1.02 (0.92, 1.12)	0.71	0.06	0.05	1.06 (0.97, 1.17)	0.21	0.09	0.05	1.10 (1.00, 1.21)	0.06
	Unspecified	-0.01	0.03	0.99 (0.94, 1.04)	0.77	-0.02	0.03	0.98 (0.93, 1.03)	0.48	0.02	0.03	1.02 (0.97, 1.07)	0.51
Treatment setting		----- reference category -----											
	Outpatient												
	Emergency	0.05	0.02	1.05 (1.01,1.09)	0.02	0.03	0.02	1.03 (0.99,1.07)	0.18	0.03	0.02	1.03 (0.99,1.07)	0.13
	Inpatient	0.04	0.02	1.04 (0.99,1.09)	0.13	0.02	0.02	1.02 (0.98,1.07)	0.37	0.02	0.02	1.02 (0.97,1.07)	0.46

1011

1012 *MD = major depression, BD = bipolar disorder, SZ = schizophrenia, OR = odds ratio, 95% CI = 95% confidence interval.*

1013 Figure 1

1014 Title: Polygenic liability for MD, BD and SZ and hazard of depression in the iPSYCH2012 cohort

1015

1016 Explanatory legend: Hazard of depression for each decile of polygenic liability relative to the bottom

1017 decile (i.e. the bottom 10% of the polygenic liability in the Danish population). Bars represent 95%


1018 confidence intervals. MD = major depressive disorder, BD = bipolar disorder, SZ = schizophrenia.

1019


1020 Figure 2


1021 Title: Polygenic liability for MD, BD and SZ and hazard of depression by age and severity

1022 Explanatory legend: (A) Hazard ratios per standard deviation (SD) increase in polygenic liabilities for
1023 MD, BD and SZ for depression diagnosis within different age ranges. (B) Hazard ratios per SD
1024 increase in polygenic liabilities for MD, BD and SZ for depression diagnosis with different ICD-10
1025 severity specifications. (C) Hazard ratios per SD increase in polygenic liabilities for MD, BD and SZ
1026 for depression diagnosis within different treatment settings. Bars represent 95% confidence intervals.
1027 MD = major depressive disorder, BD = bipolar disorder, SZ = schizophrenia.



(A)





(C)

